

Dr. Desmond J. Soares

MB.BS. F.R.A.C.S

ORTHOPAEDIC SURGEON
Internet: www.drdes.com.au

Fax: (07) 3208 7775

Shop 10, Springwood Plaza
3-15 Dennis Road
Springwood Qld 4127
Ph: (07) 3208 5552

Provider No. 37311BB
ABN 96 752 622 146

INSTRUCTIONS FOR PATIENTS FOLLOWING KNEE REPLACEMENT SURGERY

Please take some PAIN KILLERS (usually 2 Panadeine forte for an adult) about 6 hours after your operation. You may need to take 2 tablets every six hours for the first few days. After the first day, cut back to 2 Panadeine or Panadol every six hours if needed. These can be purchased over the counter.

If your pain is not settling please go to the Emergency Department of Greenslopes Hospital or your nearest public hospital. You will be checked by a doctor there who will then contact Dr. Soares if necessary.

Dr. Soares has provided you with a script for XARELTO. This is a drug to reduce the risk of developing deep vein thrombosis (DVT) and pulmonary embolism (blood clots that can travel from your leg to your lungs). It is recommended that you have a dose of XARELTO once each day for 2 weeks. Please notify Dr. Soares if you have any ooze from your wound.

Please walk at least twice a day. The only exercise you need is to walk. Work on getting your knee fully straight and bending to at least right angles (sitting on a chair with leg straight up and down). Please use crutches for the first few weeks until you feel confident. You can then migrate to a walking stick. You should use some form of walking aid for at least the first 4 weeks.

For the first 6 weeks after your operation please DO NOT drive a car.

Please leave your dressings intact until for at least 3 weeks after surgery. This will reduce the risk of developing a wound infection.

FOLLOW-UP APPOINTMENT: Your follow-up appointment will be at

[X] SHOP 10, SPRINGWOOD PLAZA, 3-15 DENNIS ROAD, SPRINGWOOD
Ph: 32085552

Please telephone once you get home, and make an appointment, so you can see Dr Soares after your operation. Please bring any X-rays you might have with you.

PLEASE GIVE TO PATIENT